

Autoxidation products of both carbohydrates and lipids are increased in uremic plasma: Is there oxidative stress in uremia?

TOSHIO MIYATA, MIN-XIN FU, KIYOSHI KUROKAWA, CHARLES VAN YPERSELE DE STRIHOU, SUZANNE R. THORPE, and JOHN W. BAYNES

Molecular and Cellular Nephrology, Institute of Medical Sciences and Department of Internal Medicine, Tokai University School of Medicine, Kanagawa, Japan; Service de Nephrologie, Universite Catholique De Louvain, Brussels, Belgium; and Department of Chemistry and Biochemistry, and School of Medicine, University of South Carolina, Columbia, South Carolina, USA

Autoxidation products of both carbohydrates and lipids are increased in uremic plasma: Is there oxidative stress in uremia?

Background. Advanced glycation end products (AGEs), formed by non-enzymatic glycation and oxidation (glycoxidation) reactions, have been implicated in the pathogenesis of several diseases, including normoglycemic uremia. AGE research in uremia has focused on the accumulation of carbohydrate-derived adducts generated by the Maillard reaction. Recent studies, however, have demonstrated that one AGE, the glycoxidation product carboxymethyllysine (CML), could be derived not only from carbohydrates but also from oxidation of polyunsaturated fatty acids *in vitro*, raising the possibility that both carbohydrate and lipid autoxidation might be increased in uremia.

Methods. To address this hypothesis, we applied gas chromatography-mass spectrometry and high performance liquid chromatography to measure protein adducts formed in uremic plasma by reactions between carbonyl compounds and protein amino groups: pentosidine derived from carbohydrate-derived carbonyls, malondialdehyde (MDA)-lysine derived from lipid-derived carbonyls, and CML originating possibly from both sources.

Results. All three adducts were elevated in uremic plasma. Plasma CML levels were mainly (>95%) albumin bound. Their levels were not correlated with fructoselysine levels and were similar in diabetic and non-diabetic patients on hemodialysis, indicating that their increase was not driven by glucose. Pentosidine and MDA-lysine were also increased in plasma to the same extent in diabetic and non-diabetic hemodialysis patients. Statistical analysis indicated that plasma levels of CML correlated weakly ($P < 0.05$) with those of pentosidine and MDA-lysine, but that pentosidine and MDA-lysine varied independently ($P > 0.5$).

Conclusions. These data suggest that the increased levels of AGEs in blood, and probably in tissues, reported in uremia implicate a broad derangement in non-enzymatic biochemistry involving alterations in autoxidation of both carbohydrates and lipids.

Key words: carboxymethyllysine, pentosidine, malondialdehyde-lysine, glycoxidation, lipoxidation, oxidative protein damage, non-enzymatic biochemistry, AGEs.

Received for publication February 17, 1998

and in revised form April 28, 1998

Accepted for publication May 1, 1998

© 1998 by the International Society of Nephrology

Proteins in the body are continuously modified by non-enzymatic glycation and oxidation (glycoxidation) reactions, also known as Maillard or browning reactions. Schiff base and Amadori adducts between protein amino groups and carbohydrate-derived carbonyls eventually yield advanced glycation end products (AGEs) [1]. AGE modified proteins increase during normal aging, but markedly increase in diabetes and uremia, damaging tissues through alterations of their structure and function [1], and stimulation of pathologic cellular responses [2–6].

The availability of sensitive and specific methods to identify some of the AGEs generated on proteins during the Maillard reaction has helped unravel the various pathways leading to the formation of AGEs in uremia. *In vitro* studies have demonstrated that pentosidine [7] originates from glucose-, ribose-, or ascorbic acid-derived carbonyl intermediates [8–11]. Similarly, N^{ϵ} -(carboxymethyl)lysine (CML) [12] is formed on proteins exposed to glucose or ascorbic acid under oxidative condition [10–13]. Both compounds have been taken as markers of carbohydrate-derived AGE adducts. More recently, however, similar *in vitro* studies have demonstrated that CML can also be derived from lipid sources such as polyunsaturated fatty acids [14], raising the possibility that AGE adducts arise from carbonyl compounds derived from both carbohydrates and lipids.

To test this hypothesis, we have measured in uremic plasma several stable protein adducts formed by reactions between carbonyl compounds and lysine amino groups in proteins [15]: pentosidine originating from carbohydrate-derived carbonyls, malondialdehyde (MDA)-lysine from lipid-derived carbonyls [16], and CML originating possibly from both carbohydrates and lipids. Our results show that there are significant increases in all three compounds, suggesting that the increased levels of AGEs in uremic patients, whether detected by immunohistochemical [11] or enzyme-linked immunosorbent assay (ELISA) techniques

Table 1. Profile of subjects examined in this study

	Group I Normal subjects (N = 9)	Group II Non-diabetic HD patients (N = 25)	Group III Diabetic HD patients (N = 21)
Age years	43.5 ± 8.1	60.6 ± 9.9 ^a	62.0 ± 9.1 ^a
Total protein g/dl	7.90 ± 0.32	7.28 ± 0.46 ^b	7.16 ± 0.48 ^a
Serum creatinine mg/dl	0.81 ± 0.11	10.73 ± 2.19 ^c	10.10 ± 2.25 ^a
Fructoselysine mmol/mol Lys	2.27 ± 0.17	2.50 ± 0.58	4.33 ± 1.14 ^c
Triglycerides mg/dl	88.8 ± 70.5	122.0 ± 91.2	121.7 ± 80.0

HD is hemodialysis.

^a $P < 0.001$ vs normal subjects

^b $P < 0.01$ vs normal subjects

^c $P < 0.001$ vs. normal and non-diabetic HD subjects

[17], result from a broad derangement in non-enzymatic biochemistry in uremia, involving alterations in autoxidation of both carbohydrates and lipids.

METHODS

Plasma samples

Plasma was obtained from 9 normal subjects with normal renal function and no proteinuria (Group I in Table 1), and 25 non-diabetic (Group II) and 21 diabetic patients undergoing hemodialysis (Group III). Characteristics of the study population are summarized in Table 1. None of the patients were on lipid lowering drugs. Informed consent was obtained from all patients. Normal renal function was defined as a serum creatinine level below 1.0 mg/dl. Proteinuria was considered to be absent if no protein was detectable in the urine with Albustix (Bayer-Sankyo, Tokyo, Japan). Hemodialysis was performed three times weekly for four hours with a dialysate containing 2 g/liter glucose, 30 mmol/liter bicarbonate, and 8 mmol/liter acetate. Plasma protein concentration was determined in fresh heparinized plasma using the Bio-Rad protein assay reagent (Richmond, CA, USA) with human albumin as a standard. Fractionation of plasma proteins was performed by gel filtration on a Sephacryl S-200 column (5.0 × 67 cm; Pharmacia, Uppsala, Sweden). Plasma triglycerides and creatinine were measured by routine methods at the central laboratory of Tokai University Hospital.

Materials

Unless otherwise indicated, reagents were of highest quality obtainable from Sigma (St. Louis, MO, USA). The preparation of fructoselysine, CML, pentosidine, and MDA-lysine standards have been described previously [14, 18, 19].

Analytical procedure

For measurement of CML and MDA-lysine, plasma samples (75 μ l) were diluted with 75 μ l 0.2 M sodium borate, pH 9.2, followed by addition of 15 μ l 5.5 M NaBH₄ in 0.1 N NaOH (final 500 mM BH₄). Reduction was carried out at 4°C overnight and then protein precipitated by addition of an equal volume of 20% trichloroacetic acid.

Protein was pelleted by centrifugation at 2000 × g for three to five minutes. The supernatant was discarded and the pellet washed with 300 μ l 10% trichloroacetic acid. Heavy labeled internal standards were added and the sample hydrolyzed in 3 ml 6 N HCl at 110°C for 24 hours. Fructoselysine was analyzed in a separate 75 μ l aliquot of plasma prepared as above except that the reduction step was omitted. The hydrolysates were dried by centrifugal evaporation (Savant Speed Vac, Farmingdale, NY, USA). Fructoselysine, CML, MDA-lysine, and lysine in plasma hydrolysates were measured as their N,O-trifluoroacetyl methyl esters by selected-ion monitoring gas chromatography/mass spectrometry (GC/MS), as described previously [14, 18, 19]. Analytes were normalized to the lysine content of the sample, following external standardization using calibration curves generated from solutions containing constant amounts of heavy labeled standards and increasing amounts of unlabeled standard.

Pentosidine was measured in hydrolysates of separate plasma samples after reduction, precipitation and acid hydrolysis as above, except that the final concentration of borohydride used for reduction was decreased to 100 mM. Hydrolysates were dried, dissolved in 10 ml water and applied to a 1 ml SP-Sephadex C-25 (Pharmacia Biotech) column equilibrated in water. The sample eluate was discarded. The column was next eluted with 20 ml 0.1 N HCl, and this eluate was also discarded. Pentosidine was eluted with 5 ml 0.1 N HCl. The eluate was dried and an aliquot then used for quantification of pentosidine by reversed-phase high performance liquid chromatography (HPLC) using post-column fluorescence detection (Ex = 328 nm, Em = 378 nm), as described previously [19]. A second aliquot of the SP-Sephadex eluate was used to measure lysine content by amino acid analysis, and pentosidine concentration was expressed relative to lysine content.

All samples assayed in this study were analyzed in single batches to exclude interassay variation. Intraassay coefficients of variation of the assays were determined by repeat analyses (N = 5) of low (normal) and high (uremic) sample pools. Coefficients of variation ranged from 5 to 15% for the various assays, where the lower number refers to the




Fig. 1. Plasma N^ε-(carboxymethyl)lysine (CML), pentosidine, and MDA-lysine levels in normal and hemodialysis subjects. CML (A), pentosidine (B), and malondialdehyde (MDA)-lysine (C) levels in acid hydrolysates of plasma from normal subjects (Group I, $N = 9$), non-diabetic hemodialysis patients (Group II, $N = 25$), and diabetic hemodialysis patients (Group III, $N = 21$), were determined by GC/MS or HPLC assay (detailed in **Methods** section). Data are expressed as means \pm SD. * $P < 0.0001$ versus normal subjects. NS is not significant.

coefficients of variation for the high standard and the higher number refers to the coefficients of variation for the low standard: pentosidine, 5 to 13%; CML, 5 to 9%; and MDA-lysine, 9 to 15%.

Statistical analysis

Data are expressed as means \pm SD. The Student's *t*-test was used for a statistical evaluation of significant difference between the two groups. Correlation was assessed by linear regression analysis.

RESULTS

As shown in Table 1, hemodialysis patients had slightly lower protein and significantly higher serum creatinine concentrations. Neither of these measures differed between the non-diabetic and diabetic hemodialysis patients. Fructoselysine was increased approximately two-fold in diabetic hemodialysis patients but was unchanged in non-diabetic hemodialysis patients, compared to controls. Triglycerides were similar in all three groups.

Results of measurement of CML, pentosidine and MDA-lysine in plasma are shown in Figure 1. Plasma CML levels (Fig. 1A) of non-diabetic (Group II in Table 1) and diabetic hemodialysis patients (Group III) were significantly higher than those of normal subjects (Group I) (0.313 ± 0.065 and 0.342 ± 0.090 vs. 0.090 ± 0.016

mmol/mol Lys, respectively, $P < 0.0001$). They were not significantly different between the non-diabetic and diabetic hemodialysis patients, a finding interpreted as evidence that CML accumulation is not driven by increased glucose concentrations. This conclusion is also supported by the lack of correlation between CML and fructoselysine (Fig. 2A). Thus, fructoselysine is a known precursor of CML *in vitro*, and its concentration was increased about twofold in the diabetic hemodialysis patients (Table 1).

Plasma pentosidine levels (Fig. 1B) of non-diabetic and diabetic hemodialysis patients were also significantly higher than those of normal subjects (11.12 ± 5.46 and 14.02 ± 7.04 vs. 1.13 ± 0.36 μ mol/mol Lys, respectively, $P < 0.0001$). They were not significantly different between non-diabetic and diabetic hemodialysis patients. Levels of pentosidine did not correlate with blood glucose or fructoselysine.

CML and pentosidine, both glycoxidation products, were significantly correlated with each other in all hemodialysis patients (Fig. 2B). In order to determine which plasma protein fraction(s) was post-translationally modified with CML, plasma was fractionated by gel filtration and the CML content was determined in the albumin and IgG fractions of a pooled plasma sample. CML values for total plasma, albumin, and IgG were 0.306, 0.576, 0.189 mmol/mol Lys, respectively. Since albumin constitutes the major




Fig. 2. (A) Correlation between plasma CML and fructoselysine in hemodialysis patients. Plasma CML levels were not correlated with those of fructoselysine in hemodialysis subjects ($N = 46$, $r = 0.186$, $P = 0.216$). The equation of the line is $y = 3.022x + 2.345$. (B) Correlation between plasma CML and pentosidine in hemodialysis patients. Plasma CML levels were correlated with those of pentosidine in hemodialysis subjects ($N = 46$, $r = 0.430$, $P < 0.01$). The equation of the line is $y = 34.86x + 1.07$.




Fig. 3. (A) Correlation between plasma pentosidine and MDA-lysine in hemodialysis patients. Plasma pentosidine levels were not correlated with those of MDA-lysine in hemodialysis subjects ($N = 46$, $r = 0.097$, $P = 0.522$). The equation of the line is $y = 0.00013x + 0.033$. (B) Correlation between plasma CML and MDA-lysine in hemodialysis patients. Plasma CML levels were correlated with those of MDA-lysine in hemodialysis subjects ($N = 46$, $r = 0.320$, $P < 0.05$). The equation of the line is $y = 0.035x + 0.023$.

protein in plasma (~40 mg albumin out of ~70 mg total plasma protein/ml), these data indicate that over 95% of plasma CML is associated with the albumin fraction, a finding similar to that previously reported for pentosidine [20].

Levels of MDA-lysine levels (Fig. 1C) in plasma proteins of non-diabetic and diabetic hemodialysis patients were also significantly higher than those of normal subjects (0.037 ± 0.009 and 0.033 ± 0.008 vs. 0.019 ± 0.005 mmol/mol Lys, respectively, $P < 0.0001$). Like CML and pentosidine, they were not significantly different between diabetic and non-diabetic hemodialysis patients. Levels of MDA-lysine were not correlated with those of triglycerides in hemodialysis patients ($r = -0.329$, $P = 0.467$).

Plasma levels of CML, but not pentosidine (Fig. 3A), were correlated with MDA levels in hemodialysis patients (Fig. 3B), although levels of CML and pentosidine were correlated weakly with one another.

DISCUSSION

Research on AGEs in uremia has focused on the accumulation of carbohydrate-derived adducts generated by the

Maillard reaction, such as, AGE-peptides [17, 21], pentosidine [20, 22, 23], and CML [24]. We have recently presented evidence that AGE generation in uremic plasma is stimulated by the accumulation of carbonyl intermediates resulting from the oxidation of glucose and ascorbic acid [11, 25]. We now demonstrate the parallel accumulation of another carbonyl compound, MDA, derived from lipid peroxidation. The accumulation in uremic circulation of carbonyl intermediates derived from various sources, such as carbohydrates, ascorbic acids, and lipids, may be described as "carbonyl stress."

The finding of elevated levels of MDA-lysine in uremic sera is of interest. Only indirect evidence on changes in MDA are available, based on measurement of thiobarbituric acid reactive substances (TBARS) whose levels were reported to be either elevated [26] or normal [27] in uremic sera. TBARS measurements are not specific for MDA as they include several other compounds, depending on assay conditions, including prostaglandin and leukotriene metabolites [28]. The present GC/MS methodology is highly sensitive and specific for MDA-lysine adducts in proteins. Unlike TBARS, it evaluates actual protein modification

resulting from lipid peroxidation. Although the polyunsaturated fatty acid composition of plasma was not measured in this study, the lack of a relationship between triglyceride levels and MDA-lysine or CML, among all subjects, suggests that increased polyunsaturated fatty acid oxidation was a major contributor to MDA-lysine formation.

The weak correlations of CML with MDA-lysine, and CML with pentosidine suggests that CML is derived not only from carbohydrate-oxidation products of the Maillard reaction, but also from lipid peroxidation products. Fu et al recently demonstrated the generation of CML from polyunsaturated fatty acids [14]. The lack of correlation between MDA-lysine and pentosidine is consistent with the observation that pentosidine originates solely from carbohydrate [8, 9] and MDA-lysine from lipid peroxidation. In addition to MDA, a wide variety of other lipid-derived carbonyl compounds might react with proteins and deserve further study [29].

AGEs such as pentosidine and CML, as well as MDA-lysine, are formed by carbonyl-amine chemistry between protein amino- and guanidino-groups and carbohydrate- or lipid-derived carbonyl compounds. The identity of the carbonyl compounds accumulating in uremia remains a matter of speculation. Several substances are probably involved as no single carbonyl precursor is known to result in the simultaneous formation of CML, pentosidine, and MDA-lysine. The lack of difference between diabetic and non-diabetic uremic plasma suggests that it is not glucose. It cannot be ascorbate/dehydroascorbate whose level is decreased in uremic plasma [11]. Another intermediate compound formed during Maillard reaction, 3-deoxyglucosone, is known to accumulate in uremia [30], but is unlikely to be a candidate. Addition of 3-deoxyglucosone to normal plasma, even at concentrations higher than those reported in uremic subjects, failed to raise pentosidine concentrations during *in vitro* incubation [25].

The present results demonstrate that, like pentosidine, over 90% of CML is bound to albumin. The two best characterized AGE adducts are thus associated with high molecular weight substances in the serum, a finding in contrast to that reported for immunologically detected AGE-peptides (molecular weight, < 10 kD) [17]. Although most, if not all polyclonal anti-AGE antibodies recognize CML [31, 32], the one used in studies of AGE-peptides apparently recognizes another as yet unknown epitope. Until the identity of the epitope, as well as the origin of these small fragments are elucidated, it will be difficult to integrate immunological observations into a broader view of carbonyl stress in uremia.

The reasons for the elevation of carbonyl compounds in uremia remain to be elucidated. It might be due to the retention of various metabolites as a result of kidney failure, that is, either defective catabolism through glomerular filtration and subsequent tubular reabsorption and

destruction as demonstrated for pentosidine [33] or impaired removal through urinary excretion.

Alternatively, a uremia-associated oxidative stress [34–39] might augment the production of carbonyl compounds as already suggested for CML generation from glyoxal or dehydroascorbate, and for pentosidine generation from arabinose or dehydroascorbate [10, 11]. The respective contributions of these two pathways require further clarification.

Whatever the identity of the carbonyl compounds and the mechanisms of their rise in uremia, we suggest that identifying the various AGE adducts as well as MDA-lysine as “carbonyl stress end products” underlies that the non-enzymatic modifications disorders observed in uremia extend beyond the borders of carbohydrate-derived Maillard reaction adducts.

ACKNOWLEDGMENTS

This study was supported by Research for the Future Program of the Japan Society for the Promotion of Science grant (96L00303) to TM, United States Public Health Service grants (NIDDK 19971) to JWB and (AG11472, P01-HL55782) to SRT, and a Juvenile Diabetes Foundation International grant to SRT.

Reprint requests to Toshio Miyata, M.D., Ph.D., Molecular and Cellular Nephrology, Institute of Medical Sciences and Department of Internal Medicine, Tokai University School of Medicine, Bohseidai, Isehara, Kanagawa 259-1193, Japan.

E-mail: t-miyata@is.icc.u-tokai.ac.jp

APPENDIX

Abbreviations used in this paper: AGEs, advanced glycation end products; CML, N^ε-(carboxymethyl)lysine; ELISA, enzyme-linked immunosorbent assay; GC/MS, gas chromatography/mass spectrometry; HPLC, high-performance liquid chromatography; MDA, malondialdehyde; TBARS, thiobarbituric acid reactive substances.

REFERENCES

1. BROWNLEE M, CERAMI A, VLISSARA H: Advanced glycosylation end products in tissue and the biochemical basis of diabetic complications. *N Engl J Med* 318:1315–1321, 1988
2. KIRSTEIN M, BRETT J, RADOFF S, OGAWA S, STERN D, VLISSARA H: Advanced protein glycosylation induces transendothelial human monocyte chemotaxis and secretion of platelet-derived growth factor: Role in vascular disease of diabetes and aging. *Proc Natl Acad Sci USA* 87:9010–9014, 1990
3. VLISSARA H, BROWNLEE M, MANOGUE KR, DINARELLO CA, PASAGIAN A: Cachectin/TNF and IL-1 induced by glucose-modified proteins: Role in normal tissue remodeling. *Science* 240:1546–1548, 1988
4. MIYATA T, INAGI R, IIDA Y, SATO M, YAMADA N, ODA O, MAEDA K, SEO H: Involvement of β 2-microglobulin modified with advanced glycation end products in the pathogenesis of hemodialysis-associated amyloidosis: Induction of human monocyte chemotaxis and macrophage secretion of tumor necrosis factor- α and interleukin 1. *J Clin Invest* 93:521–528, 1994
5. MIYATA T, HORI O, ZHANG JH, YAN SD, FERRAN L, IIDA Y, SCHMIDT AM: The receptor for advanced glycation endproducts mediates the interaction of AGE- β 2-microglobulin with human mononuclear phagocytes via an oxidant-sensitive pathway: Implication for the pathogenesis of dialysis-related amyloidosis. *J Clin Invest* 98:1088–1094, 1996
6. MIYATA T, NOTOYA K, YOSHIDA K, HORIE K, KUROKAWA K, MAEDA K, TAKETOMI S: Advanced glycation end products enhance osteoclast-induced bone resorption in cultured mouse unfractionated bone cells

- and in rats implanted subcutaneously with devitalized bone particles. *J Am Soc Nephrol* 8:260–270, 1997
7. SELL DR, MONNIER VM: Structure elucidation of a senescence cross-link from human extracellular matrix. *J Biol Chem* 264:21597–21602, 1989
 8. GRANDHEE SK, MONNIER VM: Mechanism of formation of the Maillard protein cross-link pentosidine. *J Biol Chem* 266:11649–11653, 1991
 9. DYER DG, BLACKLEDGE JA, THORPE SR, BAYNES JW: Formation of pentosidine during nonenzymatic browning of protein by glucose: Identification of glucose and other carbohydrates as possible precursors of pentosidine in vivo. *J Biol Chem* 266:11654–11660, 1991
 10. WELLS-KNECHT KJ, ZYZAK DV, LITCHFIELD JE, THORPE SR, BAYNES JW: Mechanism of autooxidative glycosylation: Identification of glyoxal and arabinose as intermediates in the autooxidative modification of protein by glucose. *Biochemistry* 34:3702–3709, 1995
 11. MIYATA T, WADA Y, CAI Z, IIDA Y, HORIE K, YASUDA Y, MAEDA K, KUROKAWA K, VAN YPERSELE DE STRIHOU C: Implication of an increased oxidative stress in the formation of advanced glycation end products in patients with end-stage renal failure. *Kidney Int* 51:1170–1181, 1997
 12. AHMED MU, THORPE SR, BAYNES JW: Identification of N^{ϵ} -carboxymethyllysine as a degradation product of fructoselysine in glycated protein. *J Biol Chem* 261:4889–4894, 1986
 13. DUNN JA, AHMED MU, MURTIASHAW MH, RICHARDSON JM, WALLA MD, THORPE SR, BAYNES JW: Reaction of ascorbate with lysine and protein under autoxidizing conditions: Formation of N^{ϵ} -(carboxymethyl)lysine by reaction between lysine and products of autoxidation of ascorbate. *Biochemistry* 29:10964–10970, 1990
 14. FU MX, REQUENA JR, JENKINS AJ, LYONS TJ, BAYNES JW, THORPE SR: The advanced glycation end product, N^{ϵ} -(carboxymethyl)lysine, is a product of lipid peroxidation and glycoxidation reactions. *J Biol Chem* 271:9982–9986, 1996
 15. MIYATA T, VAN YPERSELE DE STRIHOU C, KUROKAWA K, BAYNES JW: Alterations in non-enzymatic biochemistry in uremia: Origin and significance of “carbonyl stress” in long-term uremic complications. *Kidney Int* (in press)
 16. ESTERBAUER H, SCHAUR RJ, ZOLLNER H: Chemistry and biochemistry of 4-hydroxynonenal, malondialdehyde and related aldehydes. *Free Rad Biol Med* 11:81–128, 1991
 17. MAKITA Z, VLASSARA H, CERAMI A, BUCALA R: Immunochemical detection of advanced glycosylation end products in vivo. *J Biol Chem* 267:5133–5138, 1992
 18. REQUENA JR, FU MX, AHMED MU, JENKINS AJ, LYONS TJ, BAYNES JW, THORPE SR: Quantification of malondialdehyde and 4-hydroxynonenal adducts to lysine residues in native and oxidized human low-density lipoprotein. *J Biol Chem* 272:317–325, 1997
 19. DYER DG, DUNN JA, THORPE SR, BAILIE KE, LYONS TJ, MCCANCE DR, BAYNES JW: Accumulation of Maillard reaction products in skin collagen in diabetes and aging. *J Clin Invest* 91:2463–2469, 1993
 20. MIYATA T, UEDA Y, SHINZATO T, IIDA Y, TANAKA S, KUROKAWA K, VAN YPERSELE DE STRIHOU C, MAEDA K: Accumulation of albumin-linked and free-form pentosidine in the circulation of uremic patients with end-stage renal failure: Renal implications in the pathophysiology of pentosidine. *J Am Soc Nephrol* 7:1198–1206, 1996
 21. MAKITA Z, RADOFF S, RAYFIELD EJ, YANG Z, SKOLNIK E, DELANEY V, FRIEDMAN EA: Advanced glycosylation end products in patients with diabetic nephropathy. *N Engl J Med* 325:836–842, 1991
 22. ODETTI P, FORGARTY J, SELL DR, MONNIER VM: Chromatographic quantitation of plasma and erythrocyte pentosidine in diabetic and uremic subjects. *Diabetes* 41:153–159, 1992
 23. MIYATA T, TANEDA S, KAWAI R, UEDA Y, HORIUCHI S, HARA M, MAEDA K, MONNIER VM: Identification of pentosidine as a native structure for advanced glycation end products in β 2-microglobulin-containing amyloid fibrils in patients with dialysis-related amyloidosis. *Proc Natl Acad Sci USA* 93:2353–2358, 1996
 24. DEGENHARDT TP, GRASS L, REDDY S, THORPE SR, DIAMANDIS EP, BAYNES JW: The serum concentration of the advanced glycation end-product N^{ϵ} -(carboxymethyl)lysine is increased in uremia. *Kidney Int* 52:1064–1067, 1997
 25. MIYATA T, UEDA Y, YAMADA Y, IZUHARA Y, SAITO A, JADOUL M, KUROKAWA K, VAN YPERSELE DE STRIHOU C: Accumulation of carbonyls accelerate the formation of pentosidine, an advanced glycation end product: Carbonyl stress in uremia. *J Am Soc Nephrol* (in press)
 26. DASGUPTA A, HUSSAIN S, AHMAD S: Increased lipid peroxidation in patients on maintenance hemodialysis. *Nephron* 60:56–59, 1992
 27. PAUL JL, SALL ND, SONI T, POIGNET JL, LIDENBAUM A, MAN NK, MOATTI N, RAICHVARG D: Lipid peroxidation abnormalities in hemodialyzed patients. *Nephron* 64:106–109, 1993
 28. JANERO DR: Malondialdehyde and thiobarbituric acid-reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury. *Free Rad Bio Med* 9:515–540, 1990
 29. ZAMORA R, ALAIZ M, HIDALGO FJ: Feedback inhibition of oxidative stress by oxidized lipid/amino acid reaction products. *Biochemistry* 36:15765–15771, 1997
 30. NIWA T, KATSUZAKI T, MOMOI T, MIYAZAKI T, OGAWA H, SAITO A, MIYAZAKI S, MAEDA K, TATEMACHI N, TAKEI Y: Modification of β 2m with advanced glycation end products as observed in dialysis-related amyloidosis by 3-DG accumulating in uremic serum. *Kidney Int* 49:861–867, 1996
 31. REDDY S, BICHLER J, WELLS-KNECHT KJ, THORPE SR, BAYNES JW: N^{ϵ} -(carboxymethyl)lysine is a dominant advanced glycation end product (AGE) antigen in tissue proteins. *Biochemistry* 34:10872–10878, 1995
 32. TAKEDA A, YASUDA T, MIYATA T, MIZUNO K, LI M, YONEYAMA S, HORIE K, MAEDA K, SOBUE G: Immunohistochemical study of advanced glycation end products in aging and Alzheimer's disease brain. *Neurosci Lett* 221:17–20, 1996
 33. MIYATA T, UEDA Y, HORIE K, NANGAKU M, TANAKA S, VAN YPERSELE DE STRIHOU C, KUROKAWA K: Renal catabolism of advanced glycation end products: The fate of pentosidine. *Kidney Int* 53:416–422, 1998
 34. MIYATA T, MAEDA K, KUROKAWA K, VAN YPERSELE DE STRIHOU C: Oxidation conspires with glycation to generate noxious advanced glycation end products in renal failure. *Nephrol Dial Transplant* 12:255–258, 1997
 35. CANESTRARI F, GALLI F, GIORGINI A, ALBERTINI MC, GALIOTTA P, PASCUCCI M, BOSSU M: Erythrocyte redox state in uremic anemia: Effects of hemodialysis and relevance of glutathione metabolism. *Acta Haematol* 91:187–193, 1994
 36. YEUNG JH: Effects of glycerol-induced acute renal failure on tissue glutathione and glutathione-dependent enzymes in the rat. *Methods Find Exp Clin Pharmacol* 13:23–28, 1991
 37. KUMANO K, YOKOTA S, GO M, SUYAMA K, SAKAI T, ERA S, SOGAMI M: Quantitative and qualitative changes of serum albumin in CAPD patients. *Adv Perit Dial* 8:127–130, 1992
 38. WITKO-SARSAT V, FRIEDLANDER M, CAPELLERE-BLANDIN C, CAPELLERE-BLANDIN C, NGUYEN-KHOA T, NGUYEN AN, ZINGRAFF J, JUNGERS P, DESCAMPS-LATSCHA B: Advanced oxidation protein products as a novel marker of oxidative stress in uremia. *Kidney Int* 49:1304–1313, 1996
 39. ODETTI P, GARIBALDI S, GURRERI G, ARAGNO I, DAPINO D, PRONZATO MA: Protein oxidation in hemodialysis and kidney transplantation. *Metabolism* 45:1319–1322, 1996